



# Clinical services

Tubal ligation

## Things to know:

### Consent

- It is important that you understand the process, any risks, and that you are sure about your choice.
- We will ask you to give your written consent to the procedure.

**! If you do not understand anything in this booklet, you have any questions, or if you feel that you're being pressured or coerced, please let us know.**

### Make a complaint or provide feedback

If you would like to provide any feedback, you can:

- talk to the person you are on the phone with
- call our National Support Centre on **1300 003 707**
- use our feedback form on the homepage of our website: **[msiaustralia.org.au](https://msiaustralia.org.au)**

### Interpreter services

Please let us know if you need a telephone interpreter to help you understand any of this information. If you need one at any time during the tubal ligation of pregnancy process, you can call

**Translating and Interpreting Service National on 131 450** and ask to be connected to MSI Australia on **1300 003 707**.

# What is tubal ligation? (female sterilisation)

Tubal ligation is a permanent form of contraception for women and people with a uterus. It is a surgical procedure, performed under general anaesthetic, whereby a small clip is placed on each fallopian tube to stop the egg from travelling from the ovary to the uterus. This blocks the fallopian tubes, preventing the egg and sperm from meeting and stopping a pregnancy from developing. An egg is still released with ovulation, but it is absorbed by the body.

## **How effective is the procedure?**

Tubal ligation is more than 99.5% effective. As the procedure is not 100% effective, failures can happen, even many years following the tubal ligation. If you do become pregnant after sterilisation, there is an increased chance of having an ectopic pregnancy (pregnancy outside the uterus).

You should be able to have sexual intercourse 1 week after a tubal ligation. It's important to remember that tubal ligation does **not** protect you against sexually transmitted infections (STIs). You should still consider using condoms as protection against STIs.

## **Alternatives to tubal ligation**

There are non-permanent (reversible) methods of contraception available as an alternative to tubal ligation.

These include:

- contraceptive implant/rod in your arm
- hormonal and copper intrauterine devices
- contraceptive injections
- oral contraceptive pill.

Vasectomy is a permanent method of sterilisation for men and people with testicles.


## **How is the procedure performed?**

Tubal ligation is performed using a laparoscopic (keyhole) technique. This means that 2–3 small incisions (cuts) are made in the abdomen (tummy): 1 through the umbilicus (belly button) and the other at the pubic hairline.

Air is used to gently inflate your abdomen and then a telescopic camera is inserted through the incision in the belly button to view your fallopian tubes.

Through the second incision, a small device is inserted, which is loaded with the Filsche clip. The clip is then placed on the fallopian tube which seals the tube. The process is then repeated on the other tube.

Once both tubes are clipped, the instruments are removed, and the incision sites are closed with dissolving stitches and waterproof dressings are placed over the wounds.



### What are the risks?

Laparoscopic procedures are common and safe procedures. According to RANZCOG (Royal Australian and New Zealand College of Obstetricians and Gynaecologists) the rate of complications in all laparoscopic procedures is less than 1%, with the risk of major complications less than 0.5%.

Potential complications can include:

- damage to the internal organs, such as bowel, bladder, ureter and damage to blood vessels
- excessive bleeding
- infection
- surgical emphysema (where the air becomes trapped in your skin)
- risks of general anaesthesia
- failed procedure.

# How to prepare for your procedure

## Using reliable contraception before your procedure

It's very important that you use a reliable form of contraception until the procedure. This is to prevent a very early pregnancy that may not be detected on a pregnancy test at this time.

**!** It is expected that you will spend between 2–3 hours at the clinic, depending on your recovery time.

## Fasting

Do not eat or drink anything (including chewing gum or lollies) for 6 hours before your appointment time.

You may have a small glass of water each hour up to 2 hours before your appointment, but no other fluid.

If you don't follow this, we may need to cancel or reschedule your procedure

For more information on how to prepare for general anaesthesia, please read the **sedation information booklet**.

## After your procedure

As you will have had a general anaesthetic, you cannot drive a motor vehicle for 24 hours after your appointment. You **must** have a responsible adult to pick you up from the clinic following your procedure. For more information on how to recover from the anaesthetic please read the **sedation information booklet**.

## **Rest**

When you get home from the clinic, rest and try not to do too much. You should be able to resume normal activities after 2–3 days. The effects of the anaesthetic may make you feel tired.

## **How to avoid infection**

Avoid the following for 1 week following your procedure:

- inserting anything into the vagina, including: sexual intercourse, tampons, fingers, menstrual cups, etc
- swimming or baths (showers are allowed).

If you experience any vaginal blood loss, please only use sanitary pads (do not use tampons as they can increase a chance of infection)

## **Signs of possible infection**

You may have an infection if you have any of the following:

- fever (temperature and chills)
- red and painful wounds in your abdomen (tummy)
- pain in your abdomen (tummy) that doesn't get better with pain killers
- an offensive and smelly discharge from your vagina.

If you are concerned at all call our free aftercare line at any time on **1300 888 022**.

## **Pain management**

You may experience some pain at the site of the incisions (cuts) as well as abdominal (tummy) discomfort and bloating for the first few days after your procedure. The air used to inflate your tummy may cause bloating and shoulder tip pain for a few days.

If you have any pain you may take over the counter painkillers such as paracetamol or ibuprofen. We will also provide prescription medication.

## **Dressings**

The dressings on your abdomen need to stay in place for **5 days** after your surgery.

Gently peel back the clear plastic dressings after 5 days, and the small, white strip dressings can stay in place until they fall off (or after 10 days). The stitches will dissolve over time – there is no need to have these removed.

You may shower as normal – make sure the wound sites/dressing are patted dry after showering to keep dry. Do not rub the wound sites as this can delay healing and/or cause pain.

## **Menstrual cycle**

Your menstrual cycle (period) should not be affected by the procedure – your period should arrive as per your normal cycle.

If your period has not arrived after 6 weeks, please see your GP for follow up.



## Aftercare

Please contact our Aftercare Service on **1300 888 022** if you experience **any** of the following:

- fever (temperature and chills)
- red and painful wounds in your abdomen (tummy)
- pain in your abdomen (tummy) that doesn't get better with pain killers
- an offensive and smelly discharge from your vagina.

# Your Healthcare Rights

MSI Australia adopts the Australian Charter of Healthcare Rights. The Charter provides a framework of client rights which allows clients, families, carers and services providing health care to work together towards a safe and high quality health system, achieving the best possible outcomes.

The Charter is guided by the following principles:

**Access**

**Safety**

**Respect**

**Partnership**

**Information**

**Privacy**

**Give feedback**



As a client of MSI Australia, you have a right to:

- be treated in a professional, courteous and caring manner
- be respected regardless of your gender, age, sexual orientation, disability, religion and cultural and linguistic diversity
- have your concerns, complaints and suggestions taken seriously.

For more information about your healthcare rights and the Charter, head to [safetyandquality.gov.au](https://safetyandquality.gov.au) or scan the QR code.





